5/12615068/1020

Directions for use
Read carefully!

B. Braun 0.9% Sodium Chloride Injection B.P.

Composition
100 ml of solution contain

Active ingredients:

Sodium Chloride	0.9 g

Excipients:
Water for injections

Theoretical osmolarity:	308 mOsm/l
Titration acidity:	< 0.3 mmol/l
pH:	4.5 - 7.0

Electrolyte concentrations:

Sodium	154 mmol/l
Chloride	154 mmol/l

Pharmaceutical form
Solution for injection

Pharmaco-therapeutic group
Vehicle solution

Indications
Solvent or diluent for compatible electrolyte concentrates or drugs.

Contraindications
None

Precautions for use and special warnings
B. Braun 0.9% Sodium Chloride Injection B.P. should only be administered with particular caution to patients with
- hypernatraemia
- hyperchloraemia

Interactions
When mixing with other medicaments, possible incompatibilities should be considered.

Dosage
The quantity to be chosen depends on the desired concentration of the medicament to be dissolved.

For the use of this solution as solvent/diluent for compatible electrolyte concentrates or medicaments, the instructions for use relating to the medicament to be added should be observed.

Method of administration
Intravenous or subcutaneous injection.

Overdose
Symptoms
Overdose of B. Braun 0.9% Sodium Chloride solution may result in hypernatraemia, hyperchloraemia, hyperhydration, hyperosmolarity of the serum, and metabolic acidosis.

Emergency treatment, antidotes
Immediate stop of administration, administration of diuretics with continuous monitoring of serum electrolytes, correction of electrolyte and acid-base imbalances.

Undesirable effects
Administration of larger amounts of the solution may lead to hypernatraemia and hyperchloraemia.

Note: Patients are advised to inform their doctor or pharmacist of any adverse reaction they experience in connection with the administration of this drug.

B|BRAUN

schwarz

Dokument = 148 x 210 mm
DIN A5
2 Seiten

BN_MY+MY___5
GIF (MPc, MPw) – folded
0.9% Sodium Chloride Injection B.P.
5/12615068/1020
Production site: Penang

Lätus

9099

Font size: 9 pt.

G 200137

Usage during pregnancy
No adverse reaction has been reported. However, risk versus benefits should be assessed before using.

Storage
The product should not be stored above the temperature stated on the label.

Shelf life
The product must not be used beyond the expiry date stated on the label.

Presentation
In plastic ampoules "Mini-Plasco®" and "Mini-Plasco® Connect" of 5 ml, 10 ml, 20 ml in box of 20's.

Instruction for use/handling
The solution is supplied in single-dose containers. Discard unused contents.

Use contents immediately after opening the container.

Only to be used if solution is clear and the container undamaged.

Mini-Plasco® Handling



Mini-Plasco® Connect Handling



B|BRAUN

Product registration holder and manufactured by:
B. Braun Medical Industries Sdn. Bhd.
11900 Bayan Lepas, Penang, Malaysia.