

NAME OF THE MEDICINAL PRODUCT

Sodium Chloride B. Braun 200 mg/ml Concentrate for Solution for Infusion

COMPOSITION

1 ml of concentrate for solution for infusion contains

Sodium chloride 200 mg

Electrolytes:

Sodium 3.4 mol/l

Chloride 3.4 mol/l

1 ampoule of 10 ml contains 2.0 g Sodium chloride

1 vial of 20 ml contains 4.0 g Sodium chloride

Excipient: Water for injection.

THERAPEUTIC INDICATIONS

Hyponatremia; hypochloremia; hypotonic hyperhydration.

CONTRAINDICATIONS

Hypernatremia, hyperchloremia.

UNDESIRABLE EFFECTS

The frequency of possible adverse reactions primarily depends on the dosage and the infusion rate.

Metabolism and nutrition disorders

Hypernatremia and hyperchloremia may occur during application.

Excessively rapid infusion of solutions with a high sodium chloride concentration may lead to acute volume overload.

Gastrointestinal diseases

Excessively rapid infusion of solutions with a high sodium chloride concentration may lead to diarrhoea.

Renal and urinary disorders

Excessively rapid infusion of solutions with a high sodium chloride concentration may lead to osmotic diuresis.

Administration site conditions

Peripheral vein administration of solutions with a high sodium chloride concentration may lead to irritation and inflammation of the vein septum.

WARNINGS

Keep out of the sight and reach of children. For single use only. Discard unused contents. Only to be used if the solution is clear and container and closure are undamaged. Intravenous use past dilution.

NOTE

Not all products are registered and approved for sale in all countries or regions. Indications of use may also vary by country and region. Please contact your country representative for product availability and information.

B. Braun Melsungen AG, 34212 Melsungen, Germany, 07/2014